

11. CELLULAR SENESCENCE

13 January 2023

In all multicellular species, progressive deterioration of cellular features eventually gives rise to organismal breakdown, essentially guaranteeing mortality beyond a particular maximum lifespan. Although some forms of environmental modification, such as restricted feeding, can prolong longevity, in no case is a multicellular soma known to be immortal. Dreams of breaking this barrier pharmacologically notwithstanding, there is simply no way to prevent the relentless accumulation of deleterious somatic mutations, as most single-cell variants cannot be selectively eliminated from a terminally differentiated tissue. Here, we focus on mortality features intrinsic to unicellular species (i.e., independent of external factors such as predators and pathogens). Although individual cells do not commonly undergo senescent deterioration within the span of single cell divisions, a different view emerges on the time scale of cell lineages.

It can be difficult to contemplate the issue of senescence in unicellular species, as the concept normally applies to specific adult individuals, the identity of which is often blurred when reproduction involves binary fission. In the case of an asymmetrically reproducing cell, as in the budding yeast *S. cerevisiae*, it is natural to call the smaller product the daughter cell, although it need not always follow that the smallest member of a pair is the more rejuvenated of the two (in the case of *S. cerevisiae*, it is). Things become more ambiguous when fission is morphologically symmetrical, as even in this case there can still be asymmetrical transmission of damage. One might then view the least damaged of the two cells following a division as the offspring, although quantifying such differences between two daughter cells is difficult. An alternative strategy is to base the age of a cell on some key inherited component, such as a specific pole of a parental cell.

When viewed in this way, quite possibly all cell lineages eventually succumb to senescent decline, even if nothing more than a consequence of a series of unfortunate stochastic events. This means that senescence is an inherent feature of life, present since the beginning of biology. There is nothing intrinsically beneficial about death, but as will be discussed below, the selective advantage of prolonging the life of individual cell lineages beyond extended periods of time may be negligible, in effect stalled by a drift barrier. What remains quite unclear is the degree to which senescence proceeds more rapidly in some unicellular species than in others.

Physiological Load

Senescence is the gradual deterioration of the self-sustaining features of cells, which inevitably leads to enhanced mortality rates with increasing age. The relevant de-

terminants include: the preservation of membrane, messenger RNA, and protein integrity; clearance of cytotoxic features such as harmful metabolites and protein aggregates; and maintenance of large cellular complexes such as ribosomes and nuclear pore complexes. All biomolecules are subject to harmful chemical and physical modifications. For example, lipid molecules can experience oxidative damage resulting from free radicals released during metabolism. The same is true of RNAs, which can lead to further downstream problems when erroneous messenger RNAs are translated. Proteins can accumulate damage in a number of ways, such as deamidation of Asn to Asp and Gln to Glu, both of which lead to a negatively charged amino acid with possible effects on three-dimensional protein structure.

The fundamental question is how the accumulation of damage over the life of a cell lineage can proceed at a low enough rate to avoid overwhelming the capacity for repair and eventual cell-lineage demise. Two potential routes for avoiding extinction by senescence at the population level are: 1) a rate of damage accumulation in an average cellular lifetime less than or equal to the rate at which such damage is diluted by cell division; and/or 2) asymmetric inheritance of parental-cell damage, such that one of the two “daughter” products of cell division acquires more damage than its parent had at birth, while the other is rejuvenated (Figure 11.1).

Error catastrophe. Under the assumption that deteriorated proteins and macromolecular complexes accumulate during cell lifetimes, Medvedev (1962) and Orgel (1963) proposed an error-catastrophe hypothesis for aging, arguing in particular that progressive decline in the accuracy of the transcription and translation machinery would lead to still further accumulation of erroneous proteins. The idea here is that because transcription and translation processes involve multiple proteins (Chapter 20), as these incur errors, the error-rate itself will increase, leading to a downward spiral of cell fitness. Some contribution to error catastrophe might result from genomic mutations (below), but the mechanisms envisioned here primarily concern phenotypic errors.

The vast majority of research on this matter has focused on the cells of multicellular species. Abnormal proteins accumulate in the cells of old organisms (Holliday and Tarrant 1972; Rothstein 1975), although it is unclear whether this is a consequence of declining accuracy of transcription and/or translation. Although ribosomes in rodents do not appear to decline in terms of translational accuracy with age, ribosome activity does decline (Mori et al. 1979; Butzow et al. 1981; Filion and Laughrea 1985). There is some suggestion that transcript errors accumulate in old *S. cerevisiae* cells, although these results were obtained in cell lines with error-prone RNA polymerase, and it also remains unclear whether the error rate of transcription itself increases (as expected under the error-catastrophe hypothesis) or whether older transcripts simply accumulate more damage while circulating within cells (Vermulst et al. 2015).

Other results yield a mixed perspective on the error-catastrophe hypothesis. For example, Krisko and Radman (2013) found that progressive oxidative damage in *E. coli* magnifies the genomic mutation rate, and an *E. coli* strain engineered to be defective in translation experienced an increase in the mutation rate, but this was apparently a consequence of the activation of an error-prone DNA repair pathway (Bacher and Schimmel 2007). By applying the drug streptomycin, Edelman and

Gallant (1977) elevated the translation-error rate by $50\times$ in the same species, and found that although the misincorporation of amino acids into proteins increased substantially, cellular error levels reached a steady state, with growth potential reaching a plateau rather than exhibiting a downward spiral. Thus, these studies demonstrate that physiological damage may not always lead to a relentless increase in genomic damage by positive feedback.

Orgel (1970) eventually realized that error catastrophe is not inevitable. As noted above, the key issue is whether the error rate grows sufficiently rapidly with the accumulation of damage to outpace the rate of removal. Letting $D(t)$ be the amount of cellular damage at time t , if λ is the baseline rate of production of new cellular damage (assumed to be constant), and δ is the fractional rate of removal of prior damage, the expected amount of damage in the next time unit is $\lambda + (1 - \delta)D(t)$ (Foundations 11.1). Assuming that λ and δ remain constant, if $0 < \delta \leq 1$, a stable equilibrium level of damage equal to λ/δ is predicted, such that the input of new damage is exactly balanced by the loss of old damage. Error catastrophe requires negative δ , which means that prior damage accelerates the rate of accumulation of future damage.

Of course, a sufficiently low equilibrium level of damage might still ensure cell-lineage demise. An additional issue is that this simple computation is performed in a deterministic framework and assumes no variation among cells. As noted below, variation in damage inheritance is the expectation, and this can yield a tail of extreme individuals with low fitness. Under some conditions, this might lead to a point of no return, i.e., a sort of ratchet effect towards lineage extinction. However, as noted in Foundations 11.1, among-individual variation can also lead to population immortality, as cells with exceptionally low damage loads are promoted at the expense of damage-laden cells.

Cellular vs. population immortality. Because all progeny inherit their initial cytoplasm from their mothers, asymmetric transmission of damage provides a powerful mechanism for avoiding population senescence – although maternal cells might progressively accumulate damage that reduces cell-division potential, at least one of their daughters might be rejuvenated (Ackermann et al. 2007; Erjavec et al. 2008; Chao 2010). Under this view, the gradual senescence that accrues in some cell lineages can be offset by rejuvenation in others (Figure 11.2; Foundations 11.1), ensuring population survival. This is no different than a population of a multicellular species remaining viable despite the senescence of old individuals.

The stalked bacterium *Caulobacter crescentus*, whose swimming progeny eventually settle down themselves, provides a clear example of such an effect. Cell division is asymmetrical in this taxon, with the sedentary maternal cells experiencing a progressive loss of cell-division potential, and near-complete loss of reproductive capacity after producing ~ 200 progeny (Ackermann et al. 2003, 2007). However, the species lives on, as the physiological decline of parental cells is offset by improved states in progeny.

In contrast, something close to an equilibrium growth rate is demonstrable with *E. coli* cell lineages grown in an apparatus within which mother cells are trapped and consecutive progeny are released. *E. coli* cells divide in a morphologically symmetrical manner, but in this so-called “mother machine,” maternal cells retain

one pole dating back to the onset of the experiment, with the opposite pole being created anew at each cell division. When treated in this way, the growth rates of individual maternal cells remain constant for up to 150 divisions. Although there does appear to be an increase in the probability of mortality per division after ~ 100 generations (Wang et al. 2010), the evidence for progressive senescence is less than compelling. As discussed further below, other observations on the growth of single *E. coli* cells reveal that the cell-division times of descendant lineages eventually converge on equilibrium values (Chao 2010; Rang et al. 2011).

Molecular and Cellular Determinants

The previous section highlighted the conditions under which some or all cells within a population will experience senescence associated with physiological decline, and also how populations can remain effectively immortal when individual cell lineages are not. Here, we more explicitly consider the physical aspects of damage inheritance observed in various unicellular organisms. As noted above, asymmetric damage inheritance provides a simple mechanism by which some sublineages of cells become rejuvenated (albeit at the expense of aging in others).

In *E. coli*, protein aggregates that accumulate with age, and are presumably harmful to cell health, are progressively moved to cell poles in an entirely passive manner (Lindner et al. 2008; Winkler et al. 2010). To see how this process can lead to lineage-specific senescence (or at least cell-to-cell heterogeneity in damage accumulation), consider an aggregate incapable of movement. If such a cellular inclusion develops at the pole of a cell, it will remain at that location in all descendants inheriting the pole (Figure 11.3). Likewise, if it appears near the midpoint of a symmetrically dividing cell, then it will also succumb to a permanent pole location in all future descendants. Finally, if the aggregate is near the $1/4$ or $3/4$ point in a rod-shaped cell, upon division at the midpoint, it will be present at the midpoint of one of the daughter cells, and hence will be sequestered at the pole of one of the granddaughters. A similar mechanism appears to result in the differential sequestration of outer membrane proteins to the older poles of cells (Bergmiller et al. 2017).

In contrast to the situation in bacteria, which generally lack significant internal structure, aggregates in eukaryotic cells can be actively transported along the cytoskeleton to destinations associated with protein-management activities, including aggregation disassembly and protein refolding by chaperones (Kawaguchi et al. 2003; Kaganovich et al. 2008; Malmgren Hill and Nyström 2016). Despite such activities, however, the few eukaryotic species to be evaluated still exhibit patterns of senescence of individual lineages.

One of the more dramatic examples involves the budding yeast, *S. cerevisiae*, which has asymmetric cell division, with the daughter (the bud) being substantially reduced in size relative to the mother. The evidence supports the long-term sequestration of damage within maternal *S. cerevisiae* cells (Aguilaniu et al. 2003), with aging cells progressively experiencing higher mortality rates and higher degrees of sterility (inability to mate) (Kaerberlein 2010; Schlissel et al. 2017) (Figure 11.4). Although one of the most well-established methods for lifespan extension in animals is caloric restriction (imposition of a nutrient-sufficient diet low in calories)

(Fontana et al. 2010), such treatment has essentially no effect on lifespan in *S. cerevisiae*, which produce an average of 25 to 30 offspring per maternal cell regardless of conditions (Huberts et al. 2014).

Numerous mechanisms for aging have been suggested for yeast, including the accumulation of extrachromosomal circular DNAs, malfunctioning mitochondria, and damaged proteins (Macara and Mili 2008; Kaeberlein 2010; Denoth-Lippuner et al. 2014; Kaya et al. 2021). The nuclei of older cells also progressively experience a genome-wide loss of nucleosomes, elevated rates of chromosomal breaks, and increases in translocations, insertions of mitochondrial DNA, and transposition of mobile elements (Hu et al. 2014). Several dozen proteins are either mother-enriched or daughter-enriched, the former being biased towards those localized to plasma membranes and vacuoles, and the latter towards those involved in bud construction and genome maintenance (Yang et al. 2015). Such asymmetric inheritance must be due to active transport and/or exclusion mechanisms, which include a diffusion barrier at the bud neck preventing the passage of large complexes (Shcheprova et al. 2008). Individual molecules of ~ 135 proteins are retained within maternal *S. cerevisiae* cells for up to 28 generations, consistent with the potential for long-term accumulation of protein damage (Thayer et al. 2014).

In contrast to *S. cerevisiae*, the fission yeast *S. pombe* has morphologically symmetrical division. Yet, there is still some asymmetric inheritance of damage, with “maternal” cells (those retaining bud scars) inheriting more damaged proteins (Erjavec et al. 2008). Despite this partitioning, senescence appears not to occur under well-nourished conditions (Spivey et al. 2017) (Figure 11.4), arising primarily under stressful conditions, where cells accumulating large aggregates suffer higher mortality rates (Coelho et al. 2013). Whether such behavior reflects the predicted behavior of the model outlined in Foundations 11.1, wherein both sublineages of daughter cells evolve a finite equilibrium cell-division time and hence retain “immortality,” remains unclear.

Finally, an issue of special interest with respect to senescence in eukaryotes is the biogenesis of complex macromolecular structures and organelles during cell division. Are pre-existing assemblies retained intact across cell divisions, with entirely new complexes being assembled from newly synthesized subcomponents as daughter cells grow? Or are such features frequently disassembled and reconstructed from mixtures of old and newly synthesized components?

For many organelles (e.g., endoplasmic reticulum, mitochondria, vacuoles, and peroxisomes) in yeast, preexisting structures and their protein constituents are symmetrically inherited, with new membrane complexes (comprised of components of similar ages) being incorporated with organelle growth. For example, old nuclear-pore complexes (consisting of dozens of proteins) are inherited as a unit in yeast and remain separate from those constructed from newly synthesized components (Shcheprova et al. 2008; Menendez-Benito et al. 2013). In post-mitotic rat brain cells, proteins in large complexes appear to remain in place for extended periods of time, with $\sim 25\%$ of proteins associated with nuclear pores remaining intact after an entire year (Toyama et al. 2013). In *C. elegans*, age-related deterioration of nuclear pore complexes leads to increasingly leaky nuclei enabling the entry of increased numbers of inappropriate cytoplasmic proteins (D’Angelo et al. 2009).

These diverse observations make clear that like multicellular organisms, unicells

accrue damage over their individual lifespans. However, unlike the somas of a multicellular species, which eventually succumb to nonremovable damage accumulation, extended lineages of unicellular species can avoid permanent senescent decline as selection promotes cells with minimal damage. As will be discussed further below, these principles also apply to genomic damage, which inevitably accumulates as permanent mutations in lineages by cell-to-cell descent. Whereas specific mutant cells typically cannot be selectively removed from mixed tissues of multicellular species, leading to maladies such as cancer, they can be purged from unicellular populations by natural selection, as the cell is the unit of selection.

Evolution of Senescence

The preceding discussion makes clear that unicellular species can experience senescence, with sublineages stochastically incurring enough consecutive generations of excess damage accumulation eventually succumbing to a physiological load. Less clear, however, is the extent to which the rate of senescent decline is molded by the forces of natural selection, as opposed to being a simple passive response to selection on other cellular features.

Evolutionary theories of senescence have been developed primarily from the standpoint of multicellular organisms, focusing on the separation of the largely quiescent germline and the soma, which represents the manifestation of nearly all gene expression. As the latter is disposed after each sexual generation, its accumulated damage is expendable, whereas the expectation is that the germline will be effectively immortal.

Starting from this premise, the evolution of senescence is thought to be molded by two mechanisms (Finch 1991; Charlesworth 2000; Arking 2006). First, the pleiotropy hypothesis embraces the idea of tradeoffs between alternative fitness traits, e.g., selection for energetic investment in early reproduction subtracting from what can be invested in survival (Williams 1957). A second hypothesis focuses on the diminishing fitness payoffs of progeny produced late in life (Medawar 1952; Hamilton 1966). Like compound interest applied to a bank account, the payoffs of which grow exponentially, progeny successfully produced at a young age enter the gene pool earlier and contribute to elevated rates of expansion of the genes contributing to such a life-history strategy. As a consequence, the efficiency of natural selection operating on age-specific reproductive and survival rates must eventually decline with increasing age. In the extreme situation in which individuals late in life produce no offspring at all, there can be no selection on survival-enhancing traits expressed only beyond that point (except in the case of cross-generational care, as in social animals). As a consequence, should they exist, deleterious late-acting mutations are expected to accumulate by a combination of mutation pressure and random genetic drift.

These two hypotheses are not mutually exclusive, as there is no reason that both mechanisms cannot be operating simultaneously. Moreover, as the number of genes influencing survival and reproductive rates must constitute nearly the entire genome, it is likely that the genetic mechanisms of senescence will be diverse among phylogenetic lineages, owing to the stochastic nature of the mutation process and

the targets hit.

Unfortunately, because comparative studies of senescence in unicellular species are nearly nonexistent, the extent to which either of these hypotheses is relevant to microbes remains unclear. They are testable, however, and evidence of genetic variation for lifespan in yeast (Kaya et al. 2021) suggests a way forward. For example, the antagonistic-pleiotropy hypothesis predicts that species with higher cell-division rates would exhibit more rapid rates of demise in sublineages of cells that progressively inherit the most damage from their parental cells. The mutation-accumulation hypothesis predicts that alleles with deleterious effects on fitness will be enriched in classes of genes whose expression is confined to old mother cells. Whether genes with the postulated age-specific expression properties actually ever exist in unicellular species is an open question, but data from animal species support this argument (Rodriguez et al. 2017; Cui et al. 2019; Cheng and Kirkpatrick 2021).

Mutational Meltdown

One of the more dramatic observations from experimental gerontology is that isolated lineages of vertebrate cells generally have finite lifespans, regardless of the population size of cells. After a series of cell divisions of apparently unimpaired growth, culture loss often becomes certain. Fibroblast cell lines (derived from connective tissue) from long-lived vertebrates are capable of more divisions than those from short-lived species, and cells taken from older individuals have diminished numbers of remaining cell divisions (Martin et al. 1970; Hayflick 1977). As these results arise in cultures containing large numbers of cells, they suggest some kind of near-deterministic mechanism of sudden breakdown in cellular integrity, e.g., programmed cell death. However, observations of single-cell lineages suggest that this is not the case – the loss of cell-division potential is not abrupt but continuous throughout the lifespan of the extended lineage (Raes and Remacle 1983; Smith and Pereira-Smith 1977; Karatza and Shall 1984; Angello and Prothero 1985) (Figure 11.5).

Although the relevance of cells having an “out-of-body” experience to matters involving unicellular species is unclear, there are also cases of single-celled organisms having finite numbers of cell divisions. Most notable is the situation in ciliates, many of which have characteristic culture lifespans. The substantial body of early research focused on these kinds of observations has been extensively reviewed (Nanney 1974; Smith-Sonneborn 1981; Bell 1988). For a number of species, extinction occurs in a predictable number of cell divisions (usually in the range of 150 to 1,500 cell divisions, depending on the isolate) with low intrastain variance. Moreover, this can be true both for lineages maintained by single-cell descent and in moderately large cultures. The results do not appear to be a consequence of deterioration of the lab environment, and nuclear transplantation experiments indicate that clonal decline is caused by nuclear rather than cytoplasmic factors (Aufderheide 1984, 1987). As in the case of fibroblast cultures, close observations of the fitness features of individual *Paramecium* cells are inconsistent with a threshold reduction in cell-division potential (Figure 11.5). Rather, there is a steady decline in cell reproductive capacity with time ($\sim 0.1\%$ per generation) (Smith-Sonneborn 1979; Takagi and

Yoshida 1980; Lynch and Gabriel 1990).

Whereas the exact mechanisms underlying such behavior are not clear, the observed patterns are consistent with some sort of cumulative genomic deterioration. As first pointed out by Muller (1964), obligately asexual lineages are subject to a ratchet-like mechanism of mutation accumulation. With a high rate of genome-wide input of deleterious mutation (Chapter 4), populations of clonally reproducing cells are expected to contain a distribution of fitness classes defined by recurrent mutation pressure and selection against highly loaded individuals. This distribution is expected to be unimodal (and under many genetic models, close to Poisson in form; Chapters 3 and 23), with only a small fraction of the population contained within the best fitness class. Each generation, the possibility exists that, by chance, all individuals in this class either leave no progeny at all or produce only progeny acquiring a new deleterious mutation. Either way, the best class will have been irreversibly lost, as in the absence of sexual reproduction, there is no way to recover the prior fitness class except in the extremely rare case of a precise back-mutation. Once the best class is lost, the previously second-best class advances to be the best in the population, but it too will eventually suffer the same fate, and so on.

The natural culmination of Muller's ratchet is extinction by a process known as the mutational meltdown (Lynch and Gabriel 1990; Lynch et al. 1993; Gabriel et al. 1994). Initially, a population will typically have excess reproductive capacity, but as mean individual fitness declines, a point will eventually be reached at which each individual is just able to replace itself on a per-generation basis. At that point, any further increase in the mutation load will lead to a reduction in population size, but with fewer individuals, natural selection is less efficient (Chapter 4). This then increases the rate at which the fitness ratchet clicks, continuously driving the population to a still smaller size, culminating in a downward spiral that leads to extinction via deleterious-mutation accumulation.

As this process can operate in populations containing many thousands of asexual individuals, yielding fairly deterministic extinction times despite the stochastic accumulation of mutations, it may provide an explanation for the above-noted results. The situation for ciliates likely relates to their unique binuclear form of genetic organization, involving a germline micronucleus and a somatic macronucleus. As discussed in Chapter 10, during macronuclear maturation, all chromosomes are duplicated to very high ploidy levels (typically, several hundreds), and it is from this genome that all gene expression occurs. Although the micronucleus replicates by mitosis during clonal propagation, macronuclear division is poorly understood, but is more along the lines of random fragmentation than symmetric mitosis.

Given this genetic system, loss of viability in a ciliate population during clonal propagation must be a consequence of deleterious changes in the macronuclear genome. However, such deterioration is unlikely to involve genomic mutations at the nucleotide level. In ciliates, point mutations involving single base-pair substitutions arise at a rate of $\sim 10^{-11}$ /nucleotide site/cell division in the micronucleus (Sung et al. 2012; Long et al. 2016), the lowest known rate of any organism. Assuming the error rate is the same in the macronucleus, then because the macronuclear genome size is $\sim 10^8$ bp, only one mutation is expected to arise per haploid genome every 1000 cell divisions. Thus, the more likely path to cellular senescence in ciliates involves the amitotic (fission-like) nature of macronuclear propagation, which can lead to ran-

dom drift in chromosome copy numbers, stoichiometric imbalance, and eventual loss of entire chromosomes. Assuming simple random sampling of chromosomes following macronuclear genome duplication, Kimura (1957) produced a stochastic model of chromosome loss that yields extinction dynamics that are reasonably consistent with the existing data. Notably, extinction can be averted entirely if pre-senescent clones are allowed to undergo sexual reproduction, which results in the development of a new macronucleus from the micronuclear template.

Mutation at the nucleotide level is also unlikely to account for the behavior of vertebrate fibroblast cultures. Although the mutation rate of such cells may be as high as 10^{-9} per bp per cell division (Lynch 2010) and the haploid genome size is of order 10^9 bp, at most 10% of the vertebrate genome is under significant selection, so it is likely that < 1 functionally relevant mutation (and maybe considerably less) arises per cell division. A more likely explanation for eventual culture demise is progressive loss of telomeres from chromosome ends. Indeed, immortalized mammalian cell lines are readily obtained from cancer cells with substantial genomic rearrangements, showing that mutation accumulation is an unlikely extinction mechanism.

Summary

- The molecular constituents of cells naturally deteriorate over time, raising the possibility of senescence in unicellular organisms analogous to the aging that occurs in the somas of multicellular species. However, the asymmetric inheritance of damage by daughter cells, whether programmed or simply stochastic, can lead to the rejuvenation of one member of the pair at the expense of the other. This then ensures indefinite population survival, as in multicellular species that discard the senescing somas of aging individuals.
- Two hypotheses have been suggested for the evolution of senescence: 1) a trade-off associated with pleiotropic effects, whereby investment in early reproduction imposes costs on future survival; and 2) diminishing payoffs resulting from the compound-interest effect of early contributions to the gene pool enhancing fitness more than progeny produced later in life. Both hypotheses were constructed with multicellular species in mind, and the extent to which they apply to unicellular species remains unexplored.
- Numerous examples exist in which single-cell lineages have well-defined finite life-spans, most notably in ciliates and vertebrate cell cultures. The mechanisms driving extinction in these cases likely involve large-scale problems with chromosomal integrity and imbalance, such as telomere and/or chromosome loss, rather than the accumulation of point mutations. In such cases, rejuvenation can be accomplished by sexual outcrossing and/or inactivation of genes involved in telomere erosion.

Foundations 11.1. The physiological damage load in a cell lineage. Over time, whether stationary or in a period of active growth, a cell can be expected to acquire some reduction in physiological capacities owing to the accumulation of damaged molecules (here, assumed to accumulate at rate λ per unit time). Prior damage might also be fractionally eliminated at rate δ , such that with a starting level of damage at birth D_0 , the amount of damage accumulated over time can be expressed with the recursion equation

$$D(t+1) = \lambda + (1 - \delta)D(t). \quad (11.1.1a)$$

Note that provided $0 < \delta \leq 1$, for large t , an equilibrium level of damage is asymptotically approached. This is obtained by setting $D(t+1) = D(t)$,

$$\widehat{D} = \frac{\lambda}{\delta}. \quad (11.1.1b)$$

A special case was considered by Chao (2010), who assumed $\delta t \ll 1$, so that damage simply accumulates linearly with time before cell division, and the above-noted equilibrium is never reached. Letting D_0 denote the damage of a particular cell at birth, its damage at the time of cell division T is then

$$D(T) \simeq D_0 + \lambda T. \quad (11.1.2)$$

Suppose that some critical cell feature has to achieve a critical value P_c prior to cell division (e.g., time to achieve a threshold amount of some key cellular component), with cellular damage detracting from the rate of accumulation of P , such that

$$\frac{dP}{dt} = 1 - D(t). \quad (11.1.3)$$

Equation 11.1.3 equates the rate of product accumulation to 1.0 in the absence of damage. The solution to Equation 11.1.3, obtained after substituting Equation 11.1.2, leads to a quadratic equation defining the time to cell division,

$$P_c = P(T) = (1 - D_0)T - (\lambda/2)T^2. \quad (11.1.4)$$

Given D_0 at birth and P_c at division (a fixed parameter), Equation 11.1.4 yields the cell-division time T as a function of λ . With no damage accumulation, $\lambda = D_0 = 0$, and Equation 11.1.4 leads to the definition $P_c = T$, i.e., P_c denotes the time required to linearly build up to the checkpoint in the absence of any damage accumulation.

The prior expressions assume cells growing deterministically, with each daughter cell sharing exactly half the damage in the maternal cell at the time of fission. However, with asymmetrical cell division, the population will be heterogeneous. Letting $a \leq 1/2$ and $1 - a \geq 1/2$ be the fractions of damage transmitted to the two offspring cells (indexed as 1 and 2, with the maternal cell denoted as 0), from Equation 11.1.2 the damage transmitted to the two newborns becomes

$$D_1 = a(D_0 + \lambda T_0), \quad (11.1.5a)$$

$$D_2 = (1 - a)(D_0 + \lambda T_0). \quad (11.1.5b)$$

Substituting these expressions for D_1 and D_2 for D_0 in Equation 11.1.4, and solving yields the times to cell division for the two daughter cells as a function of the features of the maternal cell (D_0, T_0),

$$T_i = \frac{1 - D_i - \sqrt{(1 - D_i)^2 - 2P_c\lambda}}{\lambda}, \quad (11.1.6)$$

where $i = 1$ or 2 .

This overall system of equations allows one to start with a given value of T_0 for a parental cell, and then obtain the cell division times for all subsequent cells (Figure 11.6). The critical issue with respect to senescence is whether T_1 and/or T_2 increase without limits. If this were true for both daughter-cell types, runaway damage and aging of the entire population would ensue. In contrast, whereas population survival in spite of some cell senescence would occur if T_2 stabilized, i.e., the least loaded daughters do not experience ever-increasing damage.

Equilibrium requires that the reduction in damage by dilution from the mother cell be balanced by the buildup of new damage in the daughter. The equilibrium conditions are obtained by substituting $D_i = D_0$ into Equations 11.1.5a,b, so that $D_i(1-a) = a\lambda T_i$. Letting $\alpha = a/(1-a)$ and writing $T_0 = \hat{T}_1, \hat{T}_2$, yields the equilibrium physiological loads,

$$\hat{D}_1 = \hat{T}_1 \lambda \alpha, \quad (11.1.7a)$$

$$\hat{D}_2 = \hat{T}_2 \lambda / \alpha. \quad (11.1.7b)$$

Finally, substituting into Equation 11.1.4 yields the equilibrium solutions,

$$\hat{T}_1 = \frac{1 - \sqrt{1 - 2P_c \lambda (1 + 2\alpha)}}{\lambda (1 + 2\alpha)}, \quad (11.1.8a)$$

$$\hat{T}_2 = \frac{1 - \sqrt{1 - 2P_c \lambda [1 + (2/\alpha)]}}{\lambda [1 + (2/\alpha)]}. \quad (11.1.8b)$$

The conditions for equilibrium are that the terms within the square roots in the previous equations be non-negative, which requires

$$\alpha \leq \frac{1 - 2P_c \lambda}{4P_c \lambda}, \quad (11.1.9a)$$

$$1/\alpha \leq \frac{1 - 2P_c \lambda}{4P_c \lambda}. \quad (11.1.9b)$$

Thus, a key for the immortality of a cell line is that the composite quantity $P_c \lambda$ not be too large. Given the definition of P_c noted above, $P_c \lambda$ can be thought of as the total amount of damage that would be built up in a cell with minimum possible division time. With symmetric division ($\alpha = 1$), the criterion for equilibrium reduces to $P_c \lambda \leq 1/6$. With asymmetric inheritance and $P_c \lambda$ sufficiently small, both \hat{T}_1 and \hat{T}_2 reach real equilibria, but with increasing levels of damage, at most one of the cell classes can reach an equilibrium. Eventually, with high enough $P_c \lambda$, neither daughter is capable of sustained reproduction, and senescence of the entire population would ensue without some intervening rejuvenation mechanism.

Some insight into the level of fitness reduction that results from damage accumulation can be acquired by noting that for the case of symmetrical division, a damage-free cell would reproduce at age P_c , doubling the population size in that interval. In contrast, in the presence of damage accumulation, cell doubling will require time \hat{T}_1 (as defined by Equation 11.1.8a with $\alpha = 1$), so that at time P_c the population would have increased by a factor of $2^{P_c/\hat{T}_1}$. Thus, relative to a damage-free cell, the reproductive rate become $(2^{P_c/\hat{T}_1})/2$. The fractional reduction in fitness, which can be viewed as the equilibrium physiological load of damage accumulation, is then

$$L = 1 - 2^{(P_c/\hat{T}_1)-1}, \quad (11.1.10a)$$

which for moderate amounts of damage, P_c/\hat{T}_1 , simplifies to

$$L \simeq \frac{1 - (P_c/\hat{T}_1)}{\sqrt{2}}. \quad (11.1.10b)$$

The damage load can be viewed as the amount of improvement in cell fitness that could be achieved by establishing mechanisms that completely eliminate damage, i.e., $\lambda = 0$, although this view ignores any bioenergetic cost of producing and maintaining damage-control mechanisms.

Literature Cited

- Ackermann, M., A. Schauerte, S. C. Stearns, and U. Jenal. 2007. Experimental evolution of aging in a bacterium. *BMC Evol. Biol.* 7: 126.
- Ackermann, M., S. C. Stearns, and U. Jenal. 2003. Senescence in a bacterium with asymmetric division. *Science* 300: 1920.
- Aguilaniu, H., L. Gustafsson, M. Rigoulet, and T. Nyström. 2003. Asymmetric inheritance of oxidatively damaged proteins during cytokinesis. *Science* 299: 1751-1753.
- Angello, J. C., and J. W. Prothero. 1985. Clonal attenuation in chick embryo fibroblasts. *Cell Tissue Kinet.* 18: 27-43.
- Arking, R. T. 2006. *The Biology of Aging*, 3rd Ed. Oxford Univ. Press, Oxford, UK.
- Aufderheide, J. 1984. Clonal aging in *Paramecium tetraurelia*. I. Absence of evidence for a cytoplasmic factor. *Mech. Ageing Dev.* 28: 57-66.
- Aufderheide, J. 1987. Clonal aging in *Paramecium tetraurelia*. II. Evidence of functional changes in the macronucleus with age. *Mech. Ageing Dev.* 37: 265-279.
- Bacher, J. M., and P. Schimmel. 2007. An editing-defective aminoacyl-tRNA synthetase is mutagenic in aging bacteria via the SOS response. *Proc. Natl. Acad. Sci. USA* 104: 1907-1912.
- Bell, G. 1988. *Sex and Death in Protozoa: the History of an Obsession*. Cambridge Univ. Press, Cambridge, UK.
- Bergmiller, T., A. M. C. Andersson, K. Tomasek, E. Balleza, D. J. Kiviet, R. Hauschild, G. Tkačik, and C. C. Guet. 2017. Biased partitioning of the multidrug efflux pump AcrAB-TolC underlies long-lived phenotypic heterogeneity. *Science* 356: 311-315.
- Butzow, J. J., M. G. McCool, and G. L. Eichhorn. 1981. Does the capacity of ribosomes to control translation fidelity change with age? *Mech. Ageing Dev.* 15: 203-216.
- Chao, L. 2010. A model for damage load and its implications for the evolution of bacterial aging. *PLoS Genet.* 6: e1001076.
- Charlesworth, B. 2000. Fisher, Medawar, Hamilton and the evolution of aging. *Genetics* 156: 927-931.
- Cheng, C., and M. Kirkpatrick. 2021. Molecular evolution and the decline of purifying selection with age. *Nat. Commun.* 12: 2657.
- Cui, R., T. Medeiros, D. Willemsen, L. N. M. Iasi, G. E. Collier, M. Graef, M. Reichard, and D. R. Valenzano. 2019. Relaxed selection limits lifespan by increasing mutation load. *Cell* 178: 385-399.
- Coelho, M., A. Dereli, A. Haese, S. Kühn, L. Malinowska, M. E. DeSantis, J. Shorter, S. Alberti, T. Gross, and I. M. Tolić-Nørrelykke. 2013. Fission yeast does not age under favorable conditions, but does so after stress. *Curr. Biol.* 23: 1844-1852.
- D'Angelo, M. A., M. Raices, S. H. Panowski, and M. W. Hetzer. 2009. Age-dependent deterioration of nuclear pore complexes causes a loss of nuclear integrity in postmitotic cells. *Cell* 136: 284-295.
- Denoth Lippuner, A., T. Julou, and Y. Barral. 2014. Budding yeast as a model organism to study the effects of age. *FEMS Microbiol. Rev.* 38: 300-325.

- Edelmann, P., and J. Gallant. 1977. On the translational error theory of aging. *Proc. Natl. Acad. Sci. USA* 74: 3396-3398.
- Erjavec, N., M. Cvijovic, E. Klipp, and T. Nyström. 2008. Selective benefits of damage partitioning in unicellular systems and its effects on aging. *Proc. Natl. Acad. Sci. USA* 105: 18764-18769.
- Filion, A. M., and M. Laughrea. 1885. Translation fidelity in the aging mammal: studies with an accurate *in vitro* system on aged rats. *Mech. Ageing Dev.* 29: 125-142.
- Finch, C. E. 1991. *Longevity, Senescence, and the Genome*. Univ. Chicago Press, Chicago, IL.
- Fontana, L., L. Partridge, and V. D. Longo. 2010. Extending healthy life span – from yeast to humans. *Science* 328: 321-326.
- Gabriel, W., M. Lynch, and R. Bürger. 1994. Muller’s ratchet and mutational meltdowns. *Evolution* 47: 1744-1757.
- Hamilton, W. D. 1966. The moulding of senescence by natural selection. *J. Theor. Biol.* 12: 12-45.
- Hayflick, L. 1977. The cellular basis for biological aging, pp. 159-188. In C. E. Finch and L. Hayflick (eds.) *Handbook of the Biology of Aging*. Van Nostrand Reinhold, New York, NY.
- Holliday, R., and G. M. Tarrant. 1972. Altered enzymes in ageing human fibroblasts. *Nature* 238: 26-30.
- Hu, Z., K. Chen, W. Li, and J. K. Tyler. 2014. Transcriptional and genomic mayhem due to aging-induced nucleosome loss in budding yeast. *Microb. Cell.* 1: 133-136.
- Huberts, D. H., J. González, S. S. Lee, A. Litsios, G. Hubmann, E. C. Wit, and M. Heinemann. 2014. Calorie restriction does not elicit a robust extension of replicative lifespan in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 111: 11727-11731.
- Kaerberlein, M. 2010. Lessons on longevity from budding yeast. *Nature* 464: 513-519.
- Kaganovich, D., R. Kopito, and J. Frydman. 2008. Misfolded proteins partition between two distinct quality control compartments. *Nature* 454: 1088-1095.
- Karatza, C., and S. Shall. 1984. The reproductive potential of normal mouse embryo fibroblasts during culture *in vitro*. *J. Cell Sci.* 66:401-409.
- Kawaguchi, Y., J. J. Kovacs, A. McLaurin, J. M. Vance, A. Ito, and T. P. Yao. 2003. The deacetylase HDAC6 regulates aggresome formation and cell viability in response to misfolded protein stress. *Cell* 115: 727-738.
- Kaya, A., C. Z. J. Phua, M. Lee, L. Wang, A. Tyshkovskiy, S. Ma, B. Barre, W. Liu, B. R. Harrison, X. Zhao, et al. 2021. Evolution of natural lifespan variation and molecular strategies of extended lifespan in yeast. *eLife* 10: e64860.
- Kimura, M. 1957. Some problems of stochastic processes in genetics. *Ann. Math. Stat.* 28: 882-901.
- Krisko, A., and M. Radman. 2013. Phenotypic and genetic consequences of protein damage. *PLoS Genet.* 9: e1003810.
- Lindner, A. B., R. Madden, A. Demarez, E. J. Stewart, and F. Taddei. 2008. Asymmetric segregation of protein aggregates is associated with cellular aging and rejuvenation. *Proc. Natl. Acad. Sci. USA* 105: 3076-3081.
- Long, H., D. J. Winter, A. Y.-C. Chang, W. Sung, S. H. Wu, M. Balboa, R. B. R. Azevedo, R. A. Cartwright, M. Lynch, and R. A. Zufall. 2016. Low base-substitution mutation rate in the

- germline genome of the ciliate *Tetrahymena thermophila*. *Genome Biol. Evol.* 8: 3629-3639.
- Lynch, M. 2010. Evolution of the mutation rate. *Trends Genetics* 26: 345-352.
- Lynch, R. Bürger, D. Butcher, and W. Gabriel. 1993. Mutational meltdowns in asexual populations. *J. Heredity* 84: 339-344.
- Lynch, M., and W. Gabriel. 1990. Mutation load and the survival of small populations. *Evolution* 44: 1725-1737.
- Macara, I. G., and S. Mili. 2008. Polarity and differential inheritance – universal attributes of life? *Cell* 135: 801-812.
- Malmgren Hill, S., and T. Nyström. 2016. Selective protein degradation ensures cellular longevity. *eLife* 5: e17185.
- Martin, G. M., C. A. Sprague, and C. J. Epstein. 1970. Replicative life-span of cultivated human cells: effects of donor's age, tissue, and genotype. *Lab Invest.* 23: 86-92.
- Medawar, P. B. 1952. *An Unsolved Problem of Biology*. H. K. Lewis, London, UK.
- Medvedev, Z. A. 1962. A hypothesis concerning the way of coding interaction between transfer RNA and messenger RNA at the later stages of protein synthesis. *Nature* 195: 38-39.
- Menendez-Benito, V., S. J. van Deventer, V. Jimenez-Garcia, M. Roy-Luzarraga, F. van Leeuwen, and J. Neefjes. 2013. Spatiotemporal analysis of organelle and macromolecular complex inheritance. *Proc. Natl. Acad. Sci. USA* 110: 175-180.
- Mori, N., D. Mizuno, and S. Goto. 1979. Conservation of ribosomal fidelity during ageing. *Mech. Ageing Dev.* 10: 379-398.
- Muller, H. J. 1964. The relation of recombination to mutational advance. *Mutat. Res.* 1: 2-9.
- Nanney, D. L. 1974. Aging and long-term temporal regulation in ciliated protozoa: a critical review. *Mech. Ageing Dev.* 3: 81-105.
- Orgel, L. E. 1963. The maintenance of the accuracy of protein synthesis and its relevance to ageing. *Proc. Natl. Acad. Sci. USA* 49: 517-521.
- Orgel, L. E. 1970. The maintenance of the accuracy of protein synthesis and its relevance to ageing: a correction. *Proc. Natl. Acad. Sci. USA* 67: 1476.
- Raes, M., and J. Renmacle. 1983. Ageing of hamster embryo fibroblasts as the result of both differentiation and stochastic mechanisms. *Exp. Gerontol.* 18:223-240.
- Rang, C. U., A. Y. Peng, and L. Chao. 2011. Temporal dynamics of bacterial aging and rejuvenation. *Curr. Biol.* 21: 1813-1816.
- Rodríguez, J. A., U. M. Marigorta, D. A. Hughes, N. Spataro, E. Bosch, and A. Navarro. 2017. Antagonistic pleiotropy and mutation accumulation influence human senescence and disease. *Nat. Ecol. Evol.* 1: 55.
- Rothstein, M. 1975. Aging and the alteration of enzymes: a review. *Mech. Ageing Dev.* 4: 325-338.
- Schlissel, G., M. K. Krzyzanowski, F. Caudron, Y. Barral, and J. Rine. 2017. Aggregation of the Whi3 protein, not loss of heterochromatin, causes sterility in old yeast cells. *Science* 355: 1184-1187.
- Shcheprova, Z., S. Baldi, S. B. Frei, G. Gonnet, and Y. Barral. 2008. A mechanism for asymmetric

- segregation of age during yeast budding. *Nature* 454: 728-734.
- Smith, J. R., and O. Pereira-Smith. 1977. Colony size distribution as a measure of age in cultured human cells. A brief note. *Mech. Ageing Dev.* 6: 283-286.
- Smith-Sonneborn, J. 1979. DNA repair and longevity assurance in *Paramecium tetraurelia*. *Science* 203: 1115-1117.
- Smith-Sonneborn, J. 1981. Genetics and aging in protozoa. *Int. Rev. Cytol.* 73: 319-354.
- Spivey, E. C., S. K. Jones, J. R. Rybarski, F. A. Saifuddin, and I. J. Finkelstein. 2017. An aging-independent replicative lifespan in a symmetrically dividing eukaryote. *eLife* 6: e20340.
- Sung, W., A. Tucker, T. G. Doak, J. Choi, W. K. Thomas, and M. Lynch. 2012. Extraordinary genome stability in the ciliate *Paramecium tetraurelia*. *Proc. Natl. Acad. Sci. USA* 109: 19339-19344.
- Takagi, Y., and M. Yoshida. 1980. Clonal death associated with the number of fissions in *Paramecium caudatum*. *J. Cell Sci.* 41: 177-191.
- Thayer, N. H., C. K. Leverich, M. P. Fitzgibbon, Z. W. Nelson, K. A. Henderson, P. R. Gafken, J. J. Hsu, and D. E. Gottschling. 2014. Identification of long-lived proteins retained in cells undergoing repeated asymmetric divisions. *Proc. Natl. Acad. Sci. USA* 111: 14019-14026.
- Toyama, B. H., J. N. Savas, S. K. Park, M. S. Harris, N. T. Ingolia, J. R. Yates, and M. W. Hetzer. 2013. Identification of long-lived proteins reveals exceptional stability of essential cellular structures. *Cell* 154: 971-982.
- Vermulst, M., A. S. Denney, M. J. Lang, C.-W. Hung, S. Moore, M. A. Moseley, J. W. Thompson, V. Madden, J. Gauer, K. J. Wolfe, et al. 2015. Transcription errors induce proteotoxic stress and shorten cellular lifespan. *Nat. Commun.* 6: 8065.
- Wang, P., L. Robert, J. Pelletier, W. L. Dang, F. Taddei, A. Wright, and S. Jun. 2010. Robust growth of *Escherichia coli*. *Curr. Biol.* 20: 1099-1103.
- Williams, G. C. 1957. Pleiotropy, natural selection and the evolution of senescence. *Evolution* 11: 398-411.
- Winkler, J., A. Seybert, L. König, S. Pruggnaller, U. Haselmann, V. Sourjik, M. Weiss, A. S. Frangakis, A. Mogk, and B. Bukau. 2010. Quantitative and spatio-temporal features of protein aggregation in *Escherichia coli* and consequences on protein quality control and cellular ageing. *EMBO J.* 29: 910-923.
- Yang, J., M. A. McCormick, J. Zheng, Z. Xie, M. Tsuchiya, S. Tsuchiyama, H. El-Samad, Q. Ouyang, M. Kaeberlein, B. K. Kennedy, et al. 2015. Systematic analysis of asymmetric partitioning of yeast proteome between mother and daughter cells reveals “aging factors” and mechanism of lifespan asymmetry. *Proc. Natl. Acad. Sci. USA* 112: 11977-11982.

Figure 11.1. Serial buildup and dilution of cellular damage. Depending on the stochastic patterns of asymmetric damage inheritance among sublineages, the entire population of cells may remain effectively immortal, while some portions of the extended pedigree experience runaway damage and go extinct. The intensity of blue denotes the physiological load carried by a cell. The top sublineage experiences regular dilution of parental-cell damage, leading to periodic rejuvenation and persistence. The bottom sublineage represents the opposite extreme – a series of consecutive generations of inheritance of excess parental cell damage, leading to cumulative physiological deterioration. Cell lineage viability is retained so long as the rate of production of rejuvenated descendants exceeds the rate of production of permanently senescing sublineages.




Figure 11.2. The distribution of damage inheritance in pedigrees of cells. **Left)** Transmission of parental damage is perfectly symmetrical, leading to invariance in the level of damage among descendent cells. In principle, the descendant cells can reach an equilibrium level of nonlethal damage if the rate of damage removal is not overwhelmed by damage accumulation each generation. **Right)** Asymmetric inheritance leads to phenotypic variance in the damage levels in populations of cells, with some nearly damage-free cells being maintained at the expense of damage-laden cells. Two extended single-cell lineages with low levels of damage (blue) are highlighted with bold connecting lines.




Figure 11.3. Passive relocation of a cellular protein aggregate (red sphere) to the poles of symmetrically dividing cells. Because the aggregate is immobile, regardless of the starting point, it will eventually be located near the poles of descendant cells, where it will then be retained indefinitely. The numbers at the ends of cells denote the ages of poles relative to the base individual. This schematic only shows the potential distribution of aggregates descendant from the single ancestral cell, but in reality, new damage aggregates will arise in each cell and be apportioned into its descendants in the same manner. Not shown is the possibility that aggregates may also dissipate over time. Modified from Lindner et al. (2008).




Figure 11.4. Individual cells of the budding yeast *S. cerevisiae* experience progressive decline in cell-division potential, and therefore an accelerated rate of mortality with age, whereas those of the fission yeast *S. pombe* experience a constant mortality rate, as revealed by the diagonal survivorship curve (on a logarithmic scale). From Spivey et al. (2017).




Figure 11.5. **Left)** Continuous decline of cell-division potential in single-cell lines of the ciliate *Paramecium caudatum* (Takagi and Yoshida 1980). **Right)** Similar plots for fibroblasts cultured from various vertebrates: human (Smith and Pereira-Smith 1977); hamster (Raes and Remacle 1973); chicken (Angello and Prothero 1985); and mouse (Karatza and Shall 1984). The lines are least-squares regressions scaled to give a time-zero fitness equal to 1.0, except in the case of hamster. From Lynch and Gabriel (1990).




Figure 11.6. **Left)** Recursion plots for the relationship between cell-division times (T) in mother and daughter cells (the x and y axes, respectively) for the case of symmetrical cell division. Diagonal dashed lines denote potential points of equilibrium, with daughter and parental-cell division times being equal to each other. Starting with a particular value of T for a mother cell, Equation 11.1.4 defines the starting damage in that cell, which when entered into Equation 11.1.5 generates the damage in the first-generation daughter cell at the time of its division; substitution of the latter into Equation 11.1.6 then leads to the cell division time of this daughter cell; and this sequence of calculations can be done anew to yield the cell-division time in the next generation. The solid line denotes the expected relationship between mother and offspring cell-division times, which must intersect the dashed line for an equilibrium to exist, and moreover must intersect from above to the left for this equilibrium to be stable (upper left panel). The solid lines with arrows denote the successive projections of cell-division times for mothers and daughters from one generation to the next – starting with a particular maternal T_0 , the offspring T_1 is determined by reading the appropriate value for the solid line off the y axis; this time value then becomes T_0 for the next generation, and so on. In the upper panel, $P_c = 2$ and $\lambda = 0.005$, so the key parameter $P_c\lambda = 0.10$, which according to Equation 11.1.9 is sufficient to allow a stable equilibrium, as designated by the intersection of the solid and dashed lines; in this case, if T starts below the equilibrium, it progressively increases until reaching this point, and vice versa, if T starts above the equilibrium. In the lower panel, the rate of damage accumulation is too high for an equilibrium to be achieved; regardless of the starting point, T eventually increases without bound as descendants progressively accumulate more and more damage. **Right)** Actual observations on the cell-division times for parent and offspring pairs of *E. coli* cells, with the solid and open points designating the times for the daughter cells with the longest and shortest division times. Although the data are quite noisy, the regression lines for both sets of points intersect the diagonal, consistent with a non-senescent population. From Chao (2010).

